



**BRAINWARE UNIVERSITY**

**Term End Examination 2022**

**Programme – B.Tech.(CSE)-AIML-2021/B.Tech.(CSE)-DS-2021/B.Tech.(CSE)-AIML-2022/B.Tech.(CSE)-DS-2022**

**Course Name – Calculus & Linear Algebra**

**Course Code - BSCM102/BSCD102**

**( Semester I )**

**Full Marks : 60**

**Time : 2:30 Hours**

[The figure in the margin indicates full marks. Candidates are required to give their answers in their own words as far as practicable.]

**Group-A**

(Multiple Choice Type Question)

1 x 15=15

1. Choose the correct alternative from the following :

- (i) Evaluate the improper integral  $\int_{-\infty}^{\infty} \frac{1}{1+x^2} dx =$ 
  - a)  $\pi$
  - b) 1
  - c)  $\frac{\pi}{2}$
  - d) None of these
- (ii) If  $f(x)$  satisfy all the conditions of Rolle's theorem in  $[a, b]$  , then  $f'(x)$  becomes zero \_\_\_\_\_ (Select the correct option)
  - a) only at one point in  $(a, b)$
  - b) at two points in  $(a, b)$
  - c) at least one point in  $(a, b)$
  - d) none of these
- (iii) Select the value of  $\Gamma\left(\frac{1}{3}\right)\Gamma\left(\frac{2}{3}\right)$  is
  - a)  $\frac{2\pi}{\sqrt{3}}$
  - b)  $\frac{3\pi}{\sqrt{2}}$
  - c)  $\frac{\pi}{\sqrt{3}}$
  - d)  $\frac{\pi}{\sqrt{2}}$
- (iv) Test the convergence of the sequence  $\left\{1, \frac{1}{3}, \frac{1}{3^2}, \dots, \frac{1}{3^n}, \dots, \infty\right\}$ 
  - a) convergent
  - b) divergent
  - c) oscillatory
  - d) none of these
- (v) Which of the following theorem can be applied to the function  $f(x) = x^3$  in the interval  $[1,3]$  ?
  - a) Rolle's Theorem
  - b) Lagrange's Mean Value Theorem
  - c) Cauchy's Mean Value Theorem
  - d) None of these
- (vi) Compute  $\int_0^{\infty} e^{-x^2} dx =$ 
  - a)  $\pi$
  - b)  $\sqrt{\pi}$
  - c)  $\frac{\sqrt{\pi}}{2}$
  - d)  $\frac{\pi}{2}$
- (vii)

For  $x > 0$ , Identify  $\lim_{x \rightarrow 0} \frac{\sin x}{\sqrt{x}} =$

- a)  $< p style="text-align: left;"> 1$   
c) 2
- b)  $< p style="text-align: left;"> 0$   
d) None of these
- (viii)  $\lim_{n \rightarrow \infty} \left[ \frac{n}{n^2} + \frac{n}{(n^2+1^2)} + \frac{n}{(n^2+2^2)} + \dots \right]$  is equal to
- a)  $-\pi/4$   
c)  $\pi/4$
- b) 0  
d)  $\pi/3$
- (ix) Select the Lagrange's form of remainder in Taylor's theorem
- a)  $\frac{h^n(1-\theta)^{(n-1)}}{(n-1)!} f^n(a + \theta h)$   
c)  $\frac{h^n}{n!} f^n(a + \theta h)$
- b)  $\frac{h^n(1-\theta)^{(n-p)}}{p(n-1)!} f^n(a + \theta h)$   
d) None of these
- (x) Estimate the sum of series  $1 + 1/2 + 1/2^2 + \dots$  is
- a) 2  
c)  $\frac{4}{3}$
- b)  $\frac{3}{2}$   
d)  $\frac{10}{9}$
- (xi) Test the convergence of the sequence  $\{x_n\}$ , where  $x_n = (-1)^{n-1}$ , is a
- a) Convergent sequence  
c) Oscillating sequence
- b) Divergent sequence  
d) None of these
- (xii) Evaluate  $\lim_{n \rightarrow \infty} \frac{2n+1}{n+1} =$
- a) 1  
c) 3
- b) 2  
d) None of these
- (xiii) Compute  $\int_0^1 \frac{1}{\sqrt{1-x^2}} dx =$
- a)  $\pi$   
c)  $\frac{\pi}{2}$
- b) 1  
d) None of these
- (xiv) If  $f(x, y) = 0$ , then evaluate  $\frac{dy}{dx} =$
- a)  $\frac{f_x}{f_y}$   
c)  $-\frac{f_x}{f_y}$
- b)  $\frac{f_y}{f_x}$   
d)  $-\frac{f_y}{f_x}$
- (xv) If  $u = \log \frac{x^2}{y}$  then evaluate  $xu_x + yu_y =$
- a) u  
c) 1
- b) 0  
d) 2u

2. Show that  $\frac{(b-a)}{\sqrt{1-a^2}} < \sin^{-1} b - \sin^{-1} a < \frac{(b-a)}{\sqrt{1-b^2}}$  if  $0 < a < b < 1$  (3)

OR  
 Show that  $\frac{(b-a)}{\sqrt{1-a^2}} < \sin^{-1} b - \sin^{-1} a < \frac{(b-a)}{\sqrt{1-b^2}}$  if  $0 < a < b < 1$  (3)

3. Illustrate that  $\frac{x}{1+x} < \log(1+x) < x$  for all  $x > 0$ . (3)

OR  
 Use mean value theorem illustrate  $0 < \frac{1}{x} \log \frac{e^x-1}{x} < 1$ , for  $x > 0$ . (3)

4. Apply Maclaurin's theorem to the function  $f(x) = (1+x)^4$  to deduce that  $(1+x)^4 = 1 + 4x + 6x^2 + 4x^3 + x^4$ . (3)

OR  
 Determine  $\int_0^\infty e^{-x^4} x^2 dx \times \int_0^\infty e^{-x^4} dx = \frac{\pi}{8\sqrt{2}}$ . (3)

5. If  $u_n = \frac{1}{1.2} + \frac{1}{2.3} + \frac{1}{3.4} + \dots + \frac{1}{n(n+1)}$ , then explain that the sequence  $\{u_n\}$  is monotonically increasing and bounded. (3)

OR  
 If  $u = \cos^{-1} \left( \frac{x+y}{\sqrt{x}+\sqrt{y}} \right)$  then conclude that  $x \frac{\partial u}{\partial x} + y \frac{\partial u}{\partial y} + \frac{1}{2} \cot u = 0$  (3)

6. Test the extrema of the following function: (3)

$$f(x, y) = x^3 + 3xy^2 - 3y^2 - 3x^2 + 4$$

OR  
 If  $u = \log r$  and  $r^2 = x^2 + y^2 + z^2$ , Express that  $r^2 \left( \frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} + \frac{\partial^2 u}{\partial z^2} \right) = 1$ . (3)

7. Conclude that the series  $\frac{1}{1^p} + \frac{1}{2^p} + \frac{1}{3^p} + \dots$  converges for  $p > 1$  and diverges for  $p \leq 1$ . (5)

8. Tell whether the vectors  $(1,1,0)$ ,  $(1,0,1)$  and  $(0,1,1)$  form a basis of  $R^3$  over the set of real numbers. (5)

OR

Identify the linear mapping  $T : R^3 \rightarrow R^3$  which maps the basis vectors  $(0,1,1)$ ,  $(1,0,1)$ ,  $(1,1,0)$  of  $R^3$  to  $(1,1,1)$ ,  $(1,1,1)$ ,  $(1,1,1)$  respectively. (5)

9. Use the Gram-Schmidt process of orthonormalisation to identify the orthonormal basis for the sub-space of  $R^4$  generated by the vectors  $(1,1,0,1)$ ,  $(1, -2,0,0)$ ,  $(1,0,-1,2)$ . (5)

OR

Identify the values of  $a, b, c$  if the matrix  $A = \begin{bmatrix} 0 & 2b & c \\ a & b & -c \\ a & -b & c \end{bmatrix}$  is orthogonal. (5)

10. Solve  $\iint_A (x^2 + y^2) dx dy$ ; where  $A$  is the area between the line  $y = x$  and  $y = -x$  from  $x = 0$  to  $x = 1$ . Verify that the change of order does not alter the value of the integration. (5)

OR

Determine whether the limit of  $f(x, y) = \frac{xy}{y^2 - x^2}$  exist when  $(x, y) \rightarrow (0,0)$ . (5)

11. If  $\begin{vmatrix} x^3 + 3x & x - 1 & x + 3 \\ x + 1 & 1 - 2x & x - 4 \\ x - 2 & x + 4 & 3x \end{vmatrix} = ax^4 + bx^3 + cx^2 + dx + e$  be an identity in  $x$  where  $a, b, c, d, e$  are constants, then calculate the value of  $e$ . (5)

OR

Conclude that every square matrix can be uniquely expressed as sum of symmetric and skew symmetric matrix. (5)

12. Evaluate  $\iiint (x + y + z + 1)^4 dx dy dz$  over the region bounded by  $x \geq 0, y \geq 0, z \geq 0, x + y + z \leq 1$ . (5)

OR

$$\text{Justify } \int_0^1 \frac{1}{\sqrt{(1-x^4)}} dx = \frac{\sqrt{\pi}}{4} \times \frac{\Gamma(\frac{1}{4})}{\Gamma(\frac{3}{4})}$$

\*\*\*\*\*