



LIBRARY
Brahmavar University
Barasat, Kolkata - POB 126

BRAINWARE UNIVERSITY

Term End Examination 2021 - 22

Programme – Bachelor of Technology in Electronics & Communication Engineering
Course Name – Linear Algebra and Differential Equations
Course Code - BSC(ECE)201
(Semester II)

Time allotted : 1 Hrs.25 Min.

Full Marks : 70

[The figure in the margin indicates full marks.]

Group-A

(Multiple Choice Type Question)

$$1 \times 70 = 70$$

Choose the correct alternative from the following :

(1)

The value of $\begin{vmatrix} 1 & \omega & \omega^2 \\ \omega^2 & 1 & \omega \\ \omega & \omega^2 & 1 \end{vmatrix}$ is .

c) 2

- (7) The value of the determinant $\begin{vmatrix} 1 & 2 & 3 \\ 2 & 3 & 4 \\ 3 & 4 & 5 \end{vmatrix}$ is

a) 1

c) 2

- (8) If $A = \begin{pmatrix} 2 & -1 \\ 1 & 3 \end{pmatrix}$, then $A^2 + 7I =$

a) 0

c) $3A$

(9)

The rank of the matrix $A = \begin{pmatrix} 2 & 1 \\ 3 & 4 \end{pmatrix}$ is

a) 2

c) 4

- (10) For what value of μ does the system of equations $x+y+z=1$; $x+2y-z=2$; $5x+7y+\mu z=4$ have a unique solution?

a) $\mu \neq 2$

c) $\mu \neq 3$

b) $\mu \neq 1$

d) $\mu \neq 4$

(11)

The value of 'a' for which rank of the matrix $\begin{pmatrix} 2 & 0 & 1 \\ 5 & a & 3 \\ 0 & 3 & 1 \end{pmatrix}$ is less than 3?

a) $3/4$

c) $3/2$

b) $3/5$

d) 1

- (12) The equation $x-y=0$ has

a) no solution

c) exactly two solutions

b) exactly one solution

d) infinite number of solutions.

(13)

The value of $\begin{vmatrix} 100 & 101 & 102 \\ 105 & 106 & 107 \\ 110 & 111 & 112 \end{vmatrix}$ is

a) 2

c) 405

b) 0

d) -1

(14)

In $\begin{vmatrix} 3 & -2 & 5 \\ -1 & 2 & -3 \\ -5 & 6 & 9 \end{vmatrix}$, the minor and co-factor of -2 are respectively

a) -24, 24

c) -24, -24

b) 24, -24

d) none of these.

(15)

If set of vectors $\{(1, 0, 0), (1, x, 1), (x, 0, 1)\}$ is linearly dependent then x is

a) 1

c) 2

b) 0

d) 3

(16)

$S = \{(x, y, 0) | x, y \in R\}$ is a subspace of R^3 , then $\dim(S)$ is

a) 2

b) 3

c) 5

d) None of these

(17)

Let α, β, γ be three vectors in a vector space V over R, where R is the set of all real numbers. $c\alpha + d\beta + e\gamma = \theta$, where θ is the zero vector in V then the value of c, d, e are respectively.

- a) 1,1,1
- c) 1,0,0

- b) 0,0,0
- d) 0,1,1

(18)

If $\{\alpha, \beta, \gamma\}$ is a basis of a vector space V, then $\{\alpha, \beta + \gamma, \gamma\}$

- a) is a basis of V
 - c) linearly independent but not a basis
- b) linearly dependent
 - d) None of these

(19) Which of the following is not a subspace of R^3 ?

- a) $\{(x, 0) : x \in R\}$
 - b) $\{(0, y) : y \in R\}$
- c) $\{(x, 1) : x \in R\}$
 - d) $\{(x, y) : x = y; x, y \in R\}$

(20) Let $T : R^3 \rightarrow R^3$ be defined by $T(x_1, x_2, x_3) = (x_1 + 1, x_2 + 1, x_3 + 1)$, $(x_1, x_2, x_3) \in R^3$, then T is a

- a) linear mapping
 - c) $T(\alpha + \beta) = T(\alpha) + T(\beta)$
- b) is not a linear mapping
 - d) None of these

(21) Let V and W be two vector spaces and $T : V \rightarrow W$ is a linear mapping and θ, θ' be the null vectors of V and W respectively, then

- a) $Ker T = \{\alpha \in V | T(\alpha) = \theta\}$
 - c) $Ker T = \{\alpha \in V | T(\alpha) = \alpha\}$
- b) $Ker T = \{\alpha \in V | T(\alpha) = \theta'\}$
 - d) None of these

(22) If S is a subspace of a vector space $(V, +, \cdot)$ over R, where R is the set of all real numbers. Then which of the following statement is false.

- a) $\alpha + \beta \in S$ whenever $\alpha, \beta \in S$
 - c) $-\alpha + \beta \in S$ whenever $\alpha, \beta \in S$
- b) $\alpha + 2\beta \in S$ whenever $\alpha, \beta \in S$
 - d) None of a, b, c is true.

(23) Let A and B be two subspaces of a vector space V, then

- a) $A \cap B$ is a subspace of V.
 - c) $A \cup B$ is a subspace of V.
- b) both $A \cap B$ and $A \cup B$ are subspaces of V.
 - d) neither $A \cap B$ nor $A \cup B$ are subspaces of V.

(24) In a vector space V over R. Let $\alpha \in V$ and $a \in R$. Then which is true?

- a) $a\alpha \in V$
 - c) $\alpha^2 \in V$
- b) $a + \alpha \in V$
 - d) $a \in V$

(25)

The value of the linear combination $2 \begin{bmatrix} 1 & 2 & 0 \\ 2 & 1 & -1 \\ -1 & 0 & 1 \end{bmatrix} + \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$ in the vector space $M_{3,3}(R)$ is?

- a) a scalar
 - c) neither a scalar nor a vector
- b) a vector
 - d) both scalar and vector

(26) Which of the following is not linear transformation?

a) $T: R^2 \rightarrow R^2 : T(x, y) = (3x - y, 2x)$

b) $T: R^3 \rightarrow R^2 : T(x, y, z) = (3x + 1, y - z)$

c) $T: R \rightarrow R^2 : T(x) = (5x, 2x)$

d) $T: R^3 \rightarrow R^2 : T(x, y, z) = (x, 0, z)$

(27) Let I be the identity transformation of the finite dimensional vector space V , then the nullity of I is

a) $\dim(V)$

b) 0

c) 1

d) $\dim(V) - 1$

LIBRARY
Brahma Ganga University
Berhampur, Odisha - 752014

(28) A linear mapping $T: V \rightarrow W$ is injective if and only if

a) T is surjective

b)

$\text{Ker } T = \{\theta\}$

c)

d) $\text{Ker } T \neq \{\theta\}$

$\text{Im } T = \{\theta\}$

(29) Let $T: R^n \rightarrow R^m$ be a linear transformation. Which one of the following statement implies that T is bijective?

a) $\text{nullity}(T) = n$

b) $\text{rank}(T) = \text{nullity}(T) = n$

c) $\text{rank}(T) + \text{nullity}(T) = n$

d) $\text{rank}(T) - \text{nullity}(T) = n$

(30)

Which of the following is the linear transformation from R^3 to R^2 ?

(i) $f \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} 4 \\ x+y \end{pmatrix}$

(ii) $g \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} xy \\ x+y \end{pmatrix}$

(iii) $h \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} z-x \\ x+y \end{pmatrix}$

a) only f

b) only g

c) only h

d) all the transformations f,g,h

(31) Which of the following subsets of R^4 ?

$B_1 = \{(1, 0, 0, 0), (1, 1, 0, 0), (1, 1, 1, 0), (1, 1, 1, 1)\}$

$B_2 = \{(1, 0, 0, 0), (1, 2, 0, 0), (1, 2, 3, 0), (1, 2, 3, 4)\}$

$B_3 = \{(1, 2, 0, 0), (0, 0, 1, 1), (2, 1, 0, 0), (-5, 5, 0, 0)\}$

a)

B_1 and B_2 but not B_3

b)

B_1, B_2 and B_3

c)

B_1 and B_3 but not B_2

d)

only B_1

(32) If $A^2 = A$, then its Eigen values are either

a) 0 or 2

b) 1 or 2

c) 0 or 1

d) Only 0

(33) If $\lambda \neq 0$ is an Eigen value of a matrix A then the matrix A^T has an Eigen value

a) λ

b) $-\lambda$

c) $\frac{1}{\lambda}$

d) Can Not be determined

(34) If A is an orthogonal Matrix then what can we say about the matrix A

- a) Singular Matrix
- b) Non-Singular Matrix
- c) Symmetric Matrix
- d) Skew-Symmetric matrix

(35) If A is an skew-symmetric matrix then which of the following be an possible Eigen value of A

- a) 1
- b) -1
- c) 0
- d) None of -1, 0, 1

(36) If 0 is an Eigen value of a matrix A then which of the following is false

- a) 0 is an Eigen value of A^{-1}
- b) 0 is an Eigen value of A^T
- c) A has no inverse matrix
- d) A can't be orthogonal

(37) If A is an orthogonal matrix then which of the following is not a possible Eigen value of A

- a) -1
- b) 0
- c) 1
- d) $\sqrt{-1}$

(38) If A is similar to the matrix B then A^{-1} is similar to the matrix

- a) A
- b) B
- c) B^{-1}
- d) A^T

(39) If η is an Eigen value of A and A similar to B then B always has an Eigen value

- a) η^3
- b) η^2
- c) η
- d) $\frac{1}{\eta}$

(40) If α is an Eigen value and v is the corresponding Eigen vector of a matrix A then which of the following is false

- a) $Av = \alpha v$
- b) $A^T v = \alpha v$
- c) $A^{-1}v = \frac{1}{\alpha}v$
- d) One of a, b, c is false

(41) If $V = \mathbb{R}^3$ be equipped with inner product $(x, y) = x_1y_1 + 2x_2y_2 + 3x_3y_3$, In this inner

product space $(V, (\cdot, \cdot))$ then the value of the inner product of $u = \begin{bmatrix} 1 \\ 1 \\ \frac{1}{\sqrt{3}} \end{bmatrix}, v = \begin{bmatrix} 0 \\ \frac{1}{2} \\ \frac{1}{\sqrt{3}} \end{bmatrix}$

- a) $\frac{2}{\sqrt{2}}$
- b) $2\sqrt{2}$
- c) 2
- d) $\frac{\sqrt{3}}{2}$

(42) If $V = \mathbb{R}^3$ be equipped with inner product $(x, y) = x_1y_1 + x_2y_2 + x_3y_3$, In this inner product space $(V, (\cdot, \cdot))$ which of the following pairs of vectors is orthonormal?

- a)
- b)

$$u = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, v = \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}$$

$$u = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, v = \begin{bmatrix} 0 \\ \frac{1}{2} \\ 0 \end{bmatrix}$$

c) $u = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, v = \begin{bmatrix} 0 \\ 1 \\ 1 \end{bmatrix}$

d) $u = \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix}, v = \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}$

(43) Any set of linearly independent vectors can be orthonormalized by the:

- a) Cramer's rule
- b) Sobolev Method
- c) Gram-Schmidt procedure
- d) Pound-Smith procedure

(44) The diagonalizing matrix is also known as:

- a) Eigen Matrix
- b) Constant Matrix
- c) Modal Matrix
- d) State Matrix

(45) If $V = \mathbb{R}^3$ be equipped with inner product $\langle x, y \rangle = x_1y_1 + x_2y_2 + x_3y_3$. Then which of the following set of vectors are linearly independent.

- a) $\{(0, 1, 0), (0, 0, 1), (-1, 0, 1)\}$
- b) $\{(0, 1, 0), (0, -1, 0), (0, 0, 1)\}$
- c) $\{(0, 1, 0), (0, 0, 1), (-1, 0, 1)\}$
- d) $\{(1, 0, 1), (0, 1, 0), (-1, 0, 1)\}$

(46) If α and β be two orthogonal vectors in a Euclidean space $(\mathbb{R}^n, \|\cdot\|)$, then which of the following relation holds.

- a) $\|\alpha + \beta\|^2 = \|\alpha\|^2 - \|\beta\|^2$
- b) $\|\alpha + \beta\|^2 = \|\alpha\|^2 + \|\beta\|^2$
- c) $\|\alpha + \beta\|^2 = 2(\|\alpha\|^2 - \|\beta\|^2)$
- d) $\|\alpha + \beta\|^2 = 2(\|\alpha\|^2 + \|\beta\|^2)$

(47) Let A be a 3×3 matrix of real numbers and A is diagonalizable then which of the following statement is true.

- a) A has 3 l.d Eigen vectors
- b) A has 3 l.i Eigen vectors
- c) A has 3 distinct Eigen values
- d) Two of a, b and c is true

(48) If λ is an Eigen value of an orthogonal matrix A then which of the following statement is false

- a) $\det(A - \lambda I) = 0$
- b) $\det(A - \frac{1}{\lambda} I) = 0$
- c) $\det(A^{-1} - \lambda I) = 0$
- d) One of a, b and c is false

(49) If λ is the only Eigen value (real or complex) of an $n \times n$ matrix A then $\det A =$

- a) λ
- b) λ^n
- c) $n\lambda$
- d) $n\lambda^{n-1}$

(50) The differential equation $(\alpha_1 x - b_1 y) dx + (\alpha_2 x - b_2 y) dy = 0$ is exact if

- a) $\alpha_1 = b_2$
- b) $b_1 = b_2$
- c) $\alpha_1 = -b_2$
- d) $\alpha_2 = -b_1$

(51) If $x^m y^n$ be the IF of the equation $(2ydx + 3xdy) + 2xy(3ydx + 4xdy) = 0$ then the value of m and n are respectively

- a) 1, 3
- b) 2, 1
- c) 2, 2
- d) 1, 2

(52) The integrating factor of $ydx - xdy - 4x^3y^2e^x dx = 0$ is

a) $\frac{1}{y}$

b) y^2

c) xy^2

d) $\frac{1}{y^2}$

(53) The general form of a first order linear equation in x is $\frac{dy}{dx} + Py = Q$ where

- a) P and Q are both functions of x
 c) P and Q are the functions of x and y , respectively

- b) P and Q are both functions of y
 d) P and Q are the functions of y and x , respectively

(54) $\frac{1}{(D^2 - 2D + 2)} \cos x =$

a) $\frac{1}{5}(-2 \sin x + \cos x)$

b) $\frac{1}{10} \cos x$

c) $\frac{1}{5}(2 \sin x + \cos x)$

d) None of these

(55) The CF of the equation $x^2 \frac{d^2y}{dx^2} - 2x \frac{dy}{dx} - 3x = 0$ is

a) $c_1 x + c_2 e^{3x}$

b) $c_1 e^x + c_2 e^{3x}$

c) $c_1 + c_2 e^{3x}$

d) None of these

(56) The integrating factor of $\cos x \frac{dy}{dx} + y \sin x = 1$ is

a) $\tan x$

b) $\cos x$

c) $\sec x$

d) $\sin x$

(57) A particular solution of $\frac{d^2y}{dx^2} + y = 0$ when $x=0, y=4; x=\frac{\pi}{2}, y=0$ is

a) $y = A \cos x$

b) $y = 5 \cos x$

c) $y = 4 \cos x + 2 \sin x$

d) $y = 4 \cos x$

(58) $\frac{1}{(D-2)(D-3)} e^{2x} =$

a) $-e^{2x}$

b) xe^{2x}

c) $-xe^{3x}$

d) $-xe^{2x}$

(59) $\frac{1}{D^2+2} x^2 e^{3x} =$

a) $\frac{1}{11} \left(x^2 - \frac{12x}{11} \right)$

b) $\frac{1}{11} \left(x^2 - \frac{12x}{11} + \frac{60}{121} \right)$

c) $\frac{1}{11} \left(x^2 - \frac{12x}{11} + \frac{50}{121} \right)$

d) None of these

(60) The Wronskian for the differential equation $\frac{d^2y}{dx^2} - 3 \frac{dy}{dx} + 2y = 9e^x$ is

a) e^{2x}

b) e^x

c) e^{3x}

d) None of these

(61) The C.F of the equation $x^2 \frac{d^2y}{dx^2} - 2x \frac{dy}{dx} = 3x$ is

a) $c_1x + c_2e^{3x}$

c) $c_1 + c_2e^{3x}$

b) $c_1e^x + c_2e^{3x}$

d) None of these

(62) The solution of the system $Dx = y$, $Dy = x$ ($D = \frac{d}{dt}$) is

a)

$$x = Ae^t + Be^{-t}, \quad y = Ae^t + 2Be^{-t}$$

c)

$$x = Ae^t + Be^{-t}, \quad y = -Ae^t - Be^{-t}$$

b)

$$y = Ae^t + Be^{-t}, \quad x = Ae^t - Be^{-t}$$

d) None of these

(63) For the simultaneous equation $\frac{dx}{dt} + y = \sin t$, $\frac{dy}{dt} + x = \cos t$, which of the following is true?

a) $x = c_1 \cos t + c_2 \sin t$

b) $x = c_1 e^t + c_2 e^{-t}$

c) $x = (c_1 + c_2 t)e^t$

d) None of these

(64) $\frac{1}{D^2 - 2D + 5}(10 \sin x) =$

a) $\sin x + \cos x$

b) $3 \sin x - \cos x$

c) $2 \sin x + \cos x$

d) $4 \sin x$

(65) The solution of $\frac{dy}{dx} + \frac{y \cos x + \sin y + y}{\sin x + x \cos y + x} = 0$ is

a) $y \sin x + (\sin y + y) x = C$

b) $y \sin x + (\sin x + x) = C$

c) $y = \sin x + y \cos y + C$

d) None of these

(66) If y^a is an integrating factor of the differential equation $2xy \frac{dy}{dx} - (3x^2 - y^2) \frac{dy}{dx} = 0$, then the value of a is

a) -4

b) 4

c) -1

d) 1

(67) If $y = 3e^{2x} + e^{-2x} - \alpha x$ is the solution of the initial value problem $\frac{d^2y}{dx^2} + \beta y = 4\alpha x$

$$y = 4 \text{ and } \frac{dy}{dx} = 1 \text{ at } x = 0, \text{ where } \alpha, \beta \in \mathbb{R}, \text{ then}$$

a) $\alpha = 3$ and $\beta = 4$

b) $\alpha = 1$ and $\beta = 2$

c) $\alpha = 3$ and $\beta = -4$

d) $\alpha = 1$ and $\beta = -2$

(68) The integrating factor of $\frac{dy}{dx} = \frac{2xy^2 + y}{x - 2y^3}$ is

a) $\frac{1}{y}$

b) $\frac{1}{y^2}$

c) y

d) y^2

(69) Eliminating arbitrary constants a and b from $z = (x^2 + a)(y^2 + b)$, the PDE is

a) $p + q = xyz$

b) $pq = xyz$

$$c) pq = 4xyz$$

$$d) pq = -4x^2y^2z^2$$

70) The general integral of $xyp - yzq = y^2 - x^2$ for an arbitrary function ϕ is

$$a) x^2 + y^2 + z^2 = \phi(xy)$$

$$b) x^2 - y^2 - z^2 = \phi(xy)$$

$$c) x^2 + y^2 + z^2 = \phi(xyz)$$

$$d) x^2 - y^2 - z^2 = \phi(yz)$$