



HEB

**Journal of Hospital Pharmacy**  
An Official Publication of Bureau for Health & Education Status Upliftment  
(Constitutionally Entitled As Health-Education, Bureau)

JOHP

## Hypertensive urgency in a chronic kidney disease patient with hypertension after instituting Rifampicin: a case report

*Amarthya sree Racha<sup>1\*</sup>, Dr. Ashwini Soumya<sup>2</sup>*

<sup>1</sup>Amarthya sree Racha, PharmD, G. Pulla Reddy College of Pharmacy, Hyderabad, Telangana, India.

<sup>2</sup>Dr. Ashwini Soumya, Clinical pharmacist, Department of Clinical Pharmacy, AIG Hospitals, Hyderabad, Telangana, India.

\***Corresponding Author:** Racha Amarthya sree, PharmD, G. Pulla Reddy College of Pharmacy, Hyderabad, Telangana, India. Pin code: 500047

**Email Id:** [serviceheb@gmail.com](mailto:serviceheb@gmail.com)

### Abstract

Chronic kidney disease (CKD) is associated with several comorbidities and these patients have an enhanced susceptibility to tuberculosis infection owing to their immunocompromised state. Rifampicin, a first-line antitubercular drug, is a potent Cyp450 enzyme inducer. It interacts with many drugs and is known to reduce bioavailability and plasma concentration, resulting in altered drug efficacy.

Herein, we report a case of hypertensive urgency in a man with Hypertension, Diabetes mellitus, and CKD due to drug-drug interactions between Tab Metoprolol succinate (25 mg once daily), Tab Nicardipine (20 mg thrice daily), and Tab Rifampicin (600 mg once daily). The patient's blood pressure attained baseline within 7 days of modifying the antihypertensive and antitubercular therapies.

**Key words:** Chronic kidney disease, Drug-drug interactions, Enzyme induction, Hypertensive urgency, Rifampicin

### Access this Article Online

Website: <http://www.journalofhospitalpharmacy.in> Quick Response Code:

Received on 24/06/2023  
Accepted on 27/06/2023 © HEB All rights reserved

