



HEB

Journal of Hospital Pharmacy

An Official Publication of Bureau for Health & Education Status Upliftment
(Constitutionally Entitled As Health-Education, Bureau)

JOHP

Is using RABIT approach for thyroidectomy justified over open approach?

Case Title

Dr Anirban Brahma Adhikary, JR, MD, Anaesthesiology, AIIMS Rishikesh

Dr. Beeraling, Ningappa Wrikar, JR, MD Anaesthesiology, AIIMS Rishkesh

Dr. Boney John, JR, MD Anaesthesiology, AIIMS, Rishikesh

Dr Tiajem Jamir, JR, MD Anaesthesiology, AIIMS, Rishikesh

Dr. Mohammed Shafiq Shajahan, SR, MD Anaesthesiology, AIIMS, Rishikesh

Email Id: serviceheb@gmail.com

Abstract

A 48-year female came with c/o swelling on the right side of the neck for 2 years which was insidious in onset. The swelling is not associated with difficulty in swallowing, breathing or speaking. On clinical examination, it was found to be a thyroid swelling with no pressure symptoms and the patient is euthyroid status. A biopsy was done and the report revealed papillary carcinoma of the thyroid. CECT neck and thorax showed a node of size 7.7*3.7 mm in the right lobe of the thyroid with necrotic lymph nodes at cervical stations II, III and IV with no retrosternal extension and no airway involvement. Patient planned for Robotic-assisted-breast-axillo insufflation thyroidectomy with right modified radical neck dissection. The case was induced with fentanyl, propofol and vecuronium.

Keywords: RABIT, CO2 insufflation, Long duration, Hypercarbia

Access this Article Online

Website: <http://www.journalofhospitalpharmacy.in> Quick Response Code:

Received on 08/04/2022
Accepted on 22/04/2022 © HEB All rights reserved

