

# NEUROPROTECTIVE POTENTIAL OF HESPERIDIN AGAINST BENZO(A)PYRENE-INDUCED CEREBRAL PALSY

Surajit Bhattacharjee<sup>a\*</sup>, Shubhadeep Chakraborty<sup>b</sup> and Ankush Sharma<sup>c</sup>

(Received 28 March 2024) (Accepted 30 August 2024)

## ABSTRACT

This study investigates the link between cigarette smoke, specifically benzo(a)pyrene (BaP), and cerebral palsy, focusing on BaP's neurotoxicity. It also examines the neuroprotective potential of hesperidin, a citrus-derived compound, by assessing its binding affinity with the  $\beta 4$  subunit of Adapter Protein Complex 4 (AP-4). Molecular docking analyses using Discovery Studio and PyRx were performed with the  $\beta 4$  subunit of AP-4 as the protein and benzo(a)pyrene and hesperidin as ligands. The results showed that hesperidin had a higher binding affinity ( $-7.2 \text{ kcal mol}^{-1}$ ) compared to BaP ( $-6.5 \text{ kcal mol}^{-1}$ ), establishing multiple interactions, including van der Waals, hydrogen, and pi-alkyl bonds. Key receptor residues for hesperidin binding were identified. These findings highlight hesperidin's potential neuroprotective role, suggesting it may help to prevent BaP-induced neurotoxicity in cerebral palsy. The study emphasizes the promise of hesperidin, encouraging further *in vivo* and clinical investigations.