



Indian Journal of Chemistry  
Vol. 64, September 2025, pp. 899-906  
DOI: 10.56042/ijc.v64i9.18243

**NISPR**  
National Institute of Science Communication and Policy Research  
सीएसआईआर-निसप्र

## Synthesis of seven-membered oxacycle containing triquinane derivatives *via* olefin metathesis

Sambasivarao Kotha\*<sup>a,b</sup>, Kunkumita Jena<sup>#b</sup>, Ramakrishna Reddy Keesari<sup>#b</sup> & Ambreen Fatma<sup>#b</sup>

<sup>a</sup> Department of Chemistry, Sunandan Divatia School of Science, SVKM's NMIMS (Deemed to be) University, Vile Parle (West), Mumbai 400 056, India

<sup>b</sup> Department of Chemistry, Indian Institute of Technology, Bombay, Powai, Mumbai 400 076, India

E-mail: [srk@chem.iitb.ac.in](mailto:srk@chem.iitb.ac.in)

*Received 25 March 2025; accepted (revised) 22 August 2025*

Seven-membered oxacycle containing triquinane derivatives have been synthesized in good yields by employing ring-closing metathesis (RCM), ring-opening metathesis (ROM), and the ring-rearrangement metathesis (RRM) as key steps. All the compounds have been characterized by NMR data. The key building blocks have been obtained starting with a commercially available dicyclopentadiene.

**Keywords:** Triquinane, RCM, 7-Membered ring, RRM, Oxacycles

Polyquinanes containing...