

c) 2

d) 0

(7) The value of the determinant $\begin{vmatrix} 1 & 2 & 3 \\ 2 & 3 & 4 \\ 3 & 4 & 5 \end{vmatrix}$ is

a) 1

b) -1

c) 2

d) 0

(8) If $A = \begin{pmatrix} 2 & -1 \\ 1 & 3 \end{pmatrix}$, then $A^2 + 7I =$

a) O

b) 2A

c) 3A

d) 5A

(9) The rank of the matrix $A = \begin{pmatrix} 2 & 1 \\ 3 & 4 \end{pmatrix}$ is

a) 2

b) 3

c) 4

d) none of these

(10) For what value of μ does the system of equations $x+y+z=1$; $x+2y-z=2$; $5x+7y+\mu z=4$ have a unique solution?

a) $\mu \neq 2$

b) $\mu \neq 1$

c) $\mu \neq 3$

d) $\mu \neq 4$

(11) The value of 'a' for which rank of the matrix $\begin{pmatrix} 2 & 0 & 1 \\ 5 & a & 3 \\ 0 & 3 & 1 \end{pmatrix}$ is less than 3?

a) 3/4

b) 3/5

c) 3/2

d) 1

(12) The equation $x-y=0$ has

a) no solution

b) exactly one solution

c) exactly two solutions

d) infinite number of solutions.

(13)

The value of $\begin{vmatrix} 100 & 101 & 102 \\ 105 & 106 & 107 \\ 110 & 111 & 112 \end{vmatrix}$ is

a) 2

b) 0

c) 405

d) -1

(14)

In $\begin{vmatrix} 3 & -2 & 5 \\ -1 & 2 & -3 \\ -5 & 6 & 9 \end{vmatrix}$, the minor and co-factor of -2 are respectively

a) -24, 24

b) 24, -24

c) -24, -24

d) none of these.

(15)

If set of vectors $\{(1, 0, 0), (1, x, 1), (x, 0, 1)\}$ is linearly dependent then x is

a) 1

b) 0

c) 2

d) 3

(16)

$S = \{(x, y, 0) \mid x, y \in \mathbb{R}\}$ is a subspace of \mathbb{R}^3 , then $\dim(S)$ is

a) 2

b) 3

c) 5

d) None of these

(17)

Let α, β, γ be three vectors in a vector space V over \mathbb{R} , where \mathbb{R} is the set of all real numbers. $c\alpha + d\beta + e\gamma = \theta$, where θ is the zero vector in V then the value of c, d, e are respectively.

a) 1,1,1

b) 0,0,0

c) 1,0,0

d) 0,1,1

(18)

If $\{\alpha, \beta, \gamma\}$ is a basis of a vector space V , then $\{\alpha, \beta + \gamma, \gamma\}$

a) is a basis of V

b) linearly dependent

c) linearly independent but not a basis

d) None of these

(19) Which of the following is not a subspace of \mathbb{R}^2 ?

a) $\{(x, 0) : x \in \mathbb{R}\}$

b) $\{(0, y) : y \in \mathbb{R}\}$

c) $\{(x, 1) : x \in \mathbb{R}\}$

d) $\{(x, y) : x = y, x, y \in \mathbb{R}\}$

(20) Let $T : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ be defined by $T(x_1, x_2, x_3) = (x_1 + 1, x_2 + 1, x_3 + 1), (x_1, x_2, x_3) \in \mathbb{R}^3$, then T is a

a) linear mapping

b) is not a linear mapping

c) $T(\alpha + \beta) = T(\alpha) + T(\beta)$

d) None of these

(21) Let V and W be two vector spaces and $T : V \rightarrow W$ is a linear mapping and θ, θ^1 be the null vectors of V and W respectively, then

a) $\text{Ker } T = \{\alpha \in V \mid T(\alpha) = \theta\}$

b) $\text{Ker } T = \{\alpha \in V \mid T(\alpha) = \theta^1\}$

c) $\text{Ker } T = \{\alpha \in V \mid T(\alpha) = \alpha\}$

d) None of these

(22) If S is a subspace of a vector space $(V, +, \cdot)$ over \mathbb{R} , where \mathbb{R} is the set of all real numbers. Then which of the following statement is false.

a) $\alpha + \beta \in S$ whenever $\alpha, \beta \in S$

b) $\alpha + 2\beta \in S$ whenever $\alpha, \beta \in S$

c) $-\alpha + \beta \in S$ whenever $\alpha, \beta \in S$

d) None of a, b, c is true.

(23) Let A and B be two subspaces of a vector space V , then

a) $A \cap B$ is a subspace of V .

b) both $A \cap B$ and $A \cup B$ are subspaces of V .

c) $A \cup B$ is a subspace of V .

d) neither $A \cap B$ nor $A \cup B$ are subspaces of V .

(24) In a vector space V over \mathbb{R} . Let $\alpha \in V$ and $a \in \mathbb{R}$. Then which is true?

a) $a\alpha \in V$

b) $\alpha + \alpha \in V$

c) $\alpha^2 \in V$

d) $\alpha \in V$

(25)

The value of the linear combination $2 \begin{bmatrix} 1 & 2 & 0 \\ 2 & 1 & -1 \\ -1 & 0 & 1 \end{bmatrix} + \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$ in the vector space

$M_{3 \times 3}(\mathbb{R})$ is?

a) a scalar

b) a vector

c) neither a scalar nor a vector

d) both scalar and vector

(26) Which of the following is not linear transformation?

a) $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2: T(x, y) = (3x - y, 2x)$

b) $T: \mathbb{R}^3 \rightarrow \mathbb{R}^2: T(x, y, z) = (3x + 1, y - z)$

c) $T: \mathbb{R} \rightarrow \mathbb{R}^2: T(x) = (5x, 2x)$

d) $T: \mathbb{R}^3 \rightarrow \mathbb{R}^2: T(x, y, z) = (x, 0, z)$

(27) Let I be the identity transformation of the finite dimensional vector space V , then the nullity of I is

a) $\dim(V)$

b) 0

c) 1

d) $\dim(V) - 1$

(28) A linear mapping $T: V \rightarrow W$ is injective if and only if

a) T is surjective

b)

$\text{Ker } T = \{\theta\}$

c) $\text{Im } T = \{\theta\}$

d) $\text{Ker } T \neq \{\theta\}$

(29) Let $T: \mathbb{R}^n \rightarrow \mathbb{R}^n$ be a linear transformation. Which one of the following statement implies that T is bijective?

a) $\text{nullity}(T) = n$

b) $\text{rank}(T) = \text{nullity}(T) = n$

c) $\text{rank}(T) + \text{nullity}(T) = n$

d) $\text{rank}(T) - \text{nullity}(T) = n$

(30)

Which of the following is the linear transformation from \mathbb{R}^3 to \mathbb{R}^2 ?

(i) $f \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} 4 \\ x + y \end{pmatrix}$

(ii) $g \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} xy \\ x + y \end{pmatrix}$

(iii) $h \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} z - x \\ x + y \end{pmatrix}$

a) only f

b) only g

c) only h

d) all the transformations f, g, h

(31) Which of the following subsets of \mathbb{R}^4 ?

$B_1 = \{(1, 0, 0, 0), (1, 1, 0, 0), (1, 1, 1, 0), (1, 1, 1, 1)\}$

$B_2 = \{(1, 0, 0, 0), (1, 2, 0, 0), (1, 2, 3, 0), (1, 2, 3, 4)\}$

$B_3 = \{(1, 2, 0, 0), (0, 0, 1, 1), (2, 1, 0, 0), (-5, 5, 0, 0)\}$

a) B_1 and B_2 but not B_3

b) B_1, B_2 and B_3

c) B_1 and B_3 but not B_2

d) only B_1

(32) If $A^2 = A$, then its Eigen values are either

a) 0 or 2

b) 1 or 2

c) 0 or 1

d) Only 0

(33) If $\lambda \neq 0$ is an Eigen value of a matrix A then the matrix A^T has an Eigen value

a) λ

b) $-\lambda$

c) $\frac{1}{\lambda}$

d) Can Not be determined

a) $\frac{1}{y}$

b) y^2

c) xy^2

d) $\frac{1}{y^2}$

(53) The general form of a first order linear equation in x is $\frac{dy}{dx} + Px = Q$ where

a) P and Q are both functions of x b) P and Q are both functions of y c) P and Q are the functions of x and y , respectivelyd) P and Q are the functions of y and x , respectively

(54) $\frac{1}{(D^2 - 2D + 2)} \cos x =$

a) $\frac{1}{5}(-2 \sin x + \cos x)$

b) $\frac{1}{10} \cos x$

c) $\frac{1}{5}(2 \sin x + \cos x)$

d) None of these

(55) The CF of the equation $x^2 \frac{d^2y}{dx^2} - 2x \frac{dy}{dx} = 3x$ is

a) $c_1x + c_2e^{3x}$

b) $c_1e^x + c_2e^{3x}$

c) $c_1 + c_2e^{3x}$

d) None of these

(56) The integrating factor of $\cos x \frac{dy}{dx} + y \sin x = 1$ is

a) $\tan x$

b) $\cos x$

c) $\sec x$

d) $\sin x$

(57) A particular solution of $\frac{d^2y}{dx^2} + y = 0$ when $x=0, y=4; x = \frac{\pi}{2}, y=0$ is

a) $y = A \cos x$

b) $y = 5 \cos x$

c) $y = 4 \cos x + 2 \sin x$

d) $y = 4 \cos x$

(58) $\frac{1}{(D-2)(D-3)} e^{2x} =$

a) $-e^{2x}$

b) xe^{2x}

c) $-xe^{3x}$

d) $-xe^{2x}$

(59) $\frac{1}{D^2 + 2} x^2 e^{3x} =$

a) $\frac{1}{11} \left(x^2 - \frac{12x}{11} \right)$

b) $\frac{1}{11} \left(x^2 - \frac{12x}{11} + \frac{60}{121} \right)$

c) $\frac{1}{11} \left(x^2 - \frac{12x}{11} + \frac{50}{121} \right)$

d) None of these

(60) The Wronskian for the differential equation $\frac{d^2y}{dx^2} - 3 \frac{dy}{dx} + 2y = 9e^x$ is

a) e^{2x}

b) e^x

c) e^{3x}

d) None of these